§ 173.255

§ 173.255 Methylene chloride.

Methylene chloride may be present in food under the following conditions:

- (a) In spice oleoresins as a residue from the extraction of spice, at a level not to exceed 30 parts per million; *Provided*, That, if residues of other chlorinated solvents are also present, the total of all residues of such solvents shall not exceed 30 parts per million.
- (b) In hops extract as a residue from the extraction of hops, at a level not to exceed 2.2 percent, *Provided*, That:
- (1) The hops extract is added to the wort before or during cooking in the manufacture of beer.
- (2) The label of the hops extract identifies the presence of the methylene chloride and provides for the use of the hops extract only as prescribed by paragraph (b)(1) of this section.
- (c) In coffee as a residue from its use as a solvent in the extraction of caffeine from green coffee beans, at a level not to exceed 10 parts per million (0.001 percent) in decaffeinated roasted coffee and in decaffeinated soluble coffee extract (instant coffee).

§ 173.270 Hexane.

Hexane may be present in the following foods under the conditions specified:

- (a) In spice oleoresins as a residue from the extraction of spice, at a level not to exceed 25 parts per million.
- (b) In hops extract as a residue from the extraction of hops, at a level not to exceed 2.2 percent by weight; *Provided*, That:
- (1) The hops extract is added to the wort before or during cooking in the manufacture of beer.
- (2) The label of the hops extract specifies the presence of the hexane and provides for the use of the hops extract only as prescribed by paragraph (b)(1) of this section.

§ 173.275 Hydrogenated sperm oil.

The food additive hydrogenated sperm oil may be safely used in accordance with the following prescribed conditions:

(a) The sperm oil is derived from rendering the fatty tissue of the sperm whale or is prepared by synthesis of fatty acids and fatty alcohols derived

from the sperm whale. The sperm oil obtained by rendering is refined. The oil is hydrogenated.

- (b) It is used alone or as a component of a release agent or lubricant in bakery pans.
- (c) The amount used does not exceed that reasonably required to accomplish the intended lubricating effect.

§ 173.280 Solvent extraction process for citric acid.

A solvent extraction process for recovery of citric acid from conventional Aspergillus niger fermentation liquor may be safely used to produce foodgrade citric acid in accordance with the following conditions:

- (a) The solvent used in the process consists of a mixture of n-octyl alcohol meeting the requirements of §172.864 of this chapter, synthetic isoparaffinic petroleum hydrocarbons meeting the requirements of §172.882 of this chapter, and tridodecyl amine.
- (b) The component substances are used solely as a solvent mixture and in a manner that does not result in formation of products not present in conventionally produced citric acid.
- (c) The citric acid so produced meets the polynuclear aromatic hydrocarbon specifications of §173.165 and the specifications of the Food Chemicals Codex. 7th ed. (2010), pp. 226-227, which is incorporated by reference. The Director of the Office of the Federal Register approves this incorporation by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. You may obtain copies from the United States Pharmacopeial Convention, Twinbrook Pkwy., Rockville, MD 20852 (Internet address http://www.usp.org). Copies may be examined at the Food and Drug Administration's Main Library, 10903 New Hampshire Ave., Bldg. 2, Third Floor, Silver Spring, MD 20993, 301-796-2039, or at the National Archives and Records Administration (NARA). For information on the availability of this material at NARA, call 202–741–6030 or go to: http:// www.archives.gov/federal-register/cfr/ibrlocations.html.
- (d) Residues of *n*-octyl alcohol and synthetic isoparaffinic petroleum hydrocarbons are removed in accordance